

---

Citation:

Smith, DR and Duckworth, L and Jones, B (2018) Energy expenditure of elite rugby league and union players during an in-season period, measured using doubly labelled water. International Journal of Sport Nutrition and Exercise Metabolism, 28 (1). ISSN 1526-484X DOI: <https://doi.org/10.1123/ijsnem.2018-0144>

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/4595/>

Document Version:

Article (Accepted Version)

---

Published abstract

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on [openaccess@leedsbeckett.ac.uk](mailto:openaccess@leedsbeckett.ac.uk) and we will investigate on a case-by-case basis.

**Energy expenditure of elite rugby league and union players during an in-season period, measured using doubly labelled water.**

**Authors:** Smith, D.R.<sup>1, 2, 3</sup>, Duckworth, L.C.<sup>1</sup> and Jones, B.<sup>1, 3, 4</sup>.

<sup>1</sup> Leeds Beckett University, Institute for Sport, Physical Activity and Leisure, Leeds, United Kingdom.

<sup>2</sup> Leeds Rhinos RLFC, Leeds, United Kingdom

<sup>3</sup> Yorkshire Carnegie RUFC, Leeds, United Kingdom

<sup>4</sup> The Rugby Football League, Leeds, United Kingdom

Total energy expenditure (TEE) has been quantified in elite senior rugby league (RL) and rugby union (RU) players using multiple measures, with criterion measures lacking in RU and academy players. Robust measures of TEE are required as prediction equations used to estimate energy requirements are often unsuitable for athletes. This study quantified TEE of 27 elite male English academy (U16 and U20) and senior (U24) RL and RU players during a 14-day in-season period using doubly labelled water (DLW). Resting metabolic rate (RMR), using indirect calorimetry, and physical activity level (PAL) was also measured (TEE:RMR). Predicted TEE, determined by published equations, was compared to measured TEE by age group. Differences in TEE (RL,  $4369 \pm 979$ ; RU,  $4365 \pm 1122$ ; U16,  $4010 \pm 744$ ; U20,  $4414 \pm 688$ ; U24,  $4761 \pm 1523$  Kcal·day<sup>-1</sup>) and PAL (overall mean  $2.0 \pm 0.4$ ) were *unclear*. RMR was *very likely* greater for RL ( $2366 \pm 296$  Kcal·day<sup>-1</sup>) than RU players ( $2123 \pm 269$  Kcal·day<sup>-1</sup>). Relative RMR for U16, U20 and U24 ( $27 \pm 4$ ,  $23 \pm 3$  and  $26 \pm 5$  Kcal.Kg<sup>-1</sup>.day<sup>-1</sup>) was *very likely* greater for U20 than U24 players. Differences in TEE estimated by the Schofield, Cunningham and Harris-Benedict equations compared with DLW were *unclear*, *likely* and

*unclear* for U16 ( $187 \pm 614$ ;  $-489 \pm 564$  and  $-90 \pm 579$  Kcal.day<sup>-1</sup>), *likely*, *very likely* and *likely* for U20 ( $-449 \pm 698$ ;  $-785 \pm 650$  and  $-452 \pm 684$  Kcal.day<sup>-1</sup>) and all *unclear* for U24 players ( $-428 \pm 1292$ ;  $-605 \pm 1493$  and  $-461 \pm 1314$  Kcal.day<sup>-1</sup>). Due to large variability between individuals, negligible differences in TEE were observed by code, and ~350-400 Kcal.day<sup>-1</sup> differences between consecutive age groups were *unclear*. Differences in RMR may be due to training exposure and match play. The remaining components of TEE (i.e. thermic effect of feeding and activity thermogenesis) may reflect the differences in contact demands between codes, as RU players typically engage in more static exertions than RL players during match play. Prediction equations are currently insufficient to differentiate between individual variability in TEE. The importance of practitioners providing individual support for the elite rugby player is highlighted. Finally, the TEE measured in this study using the gold standard DLW method can be used as reference data for elite rugby players of different codes and ages, during an in-season training period.